



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

On February 7, with the temperature at 15 above zero, a flock of five Canada Geese passed over the city and alighted in East Potomac Park. The wind on that day at times attained a velocity of forty-five miles an hour.—BRENT M. MORGAN, 224 Eleventh St., S. W., Washington, D. C.

**Nesting of the Greater Yellow-Legs in Newfoundland.**—On June 20, 1919, Mr. J. R. Whitaker and the writer had the satisfaction of discovering a female of this species (*Totanus melanoleucus*) brooding four young just out of the shell and still in the nest, in a large bog in the vicinity of Grand Lake, N. F. Led to the spot by the ever increasing cries of the male bird, the nest, which was nothing more than a bare depression ten inches in diameter and three inches deep, upon the top of a mound of peat otherwise covered over with a short growth of sheep laurel, was noticed three yards from where we had stopped in doubt as to where next to proceed.

It presented an unusual domestic picture; one youngster was perched on the mother's back, while one or two others appeared from under her wings after the manner of domestic fowls. The parent remained until we closed in, when she flew low from the nest with a piercing cry, and after circling about overhead took up a position on a dead stub nearby, from which she continued to *kip, kip, kip, kip*—incessantly as long as we remained near the nest, the male likewise calling and circling above.

The young, whose legs were not as yet strong enough to bear their weight, lay flat in the nest. They were mottled in gray, brown and black down, white below. Some of the lighter spaces on the back tending toward buffy. The eyes were large and black, bill one-half an inch long, lead-black in color, while the legs were characteristically long and greenish in color. Notwithstanding the recent hatching of the eggs, only one or two small pieces were to be found, the empty shells doubtless having been carried away by the parents.

On visiting the nest the day following, the young could not be found, although the actions of the old birds indicated their presence in the vicinity.—GEORGE H. STUART, 3rd, Girard Trust Co., Philadelphia.

**Nesting of the Little Black Rail in Atlantic County, N. J.**—On July 4, 1919, Mr. Julian K. Potter and the writer flushed a small rail in a marsh an acre or two in extent, beyond the sand dunes immediately back of the ocean beach, on an island below Beach Haven, N. J. Searching for the nest in the belief that the bird was a Little Black Rail, we were rewarded by finding it placed among the long grasses, the tops of which were so drawn over as to almost completely hide the eggs from view. The nest, which was composed entirely of the same rather fine grass, was placed about one inch above the damp ground and contained eight eggs, very heavily incubated.